

Discrete Mathematics

Counting

Dr. Mohammad Salah Uddin
Department of Computer Science and Engineering
East West University, Bangladesh

Counting

- Assume we have a set of **objects with certain properties**
- **Counting** is used to determine **the number of these objects**

Examples:

- Number of available phone numbers with 7 digits in the local calling area

Basic counting rules

- Counting problems may be very hard, not obvious
- **Solution:**
 - **simplify the solution by decomposing the problem**
- **Two basic decomposition rules:**
 - **Product rule**
 - A count decomposes into a sequence of dependent counts (“each element in the first count is associated with all elements of the second count”)
 - **Sum rule**
 - A count decomposes into a set of independent counts (“elements of counts are alternatives”)

Product rule

A count can be broken down into a sequence of dependent counts

- “each element in the first count is associated with all elements of the second count”

Example:

- Assume an auditorium with a seat labeled by a letter and numbers in between 1 to 50 (e.g. A23). We want the total number of seats in the auditorium.
- 26 letters and 50 numbers
- How to count?
- **One solution: write down all seats (objects) and count them**

A-1 A-2 A-3 ...A-50 B-1... Z-49 Z-50

1 2 3 50 51 ... (n-1) n ← eventually we get it

Product rule

A count can be broken down into a sequence of dependent counts

- “each element in the first count is associated with all elements of the second count”

Example:

- Assume an auditorium with a seat labeled by a letter and numbers in between 1 to 50 (e.g. A23). We want the total number of seats in the auditorium.
- 26 letters and 50 numbers

- 26 letters and 50 numbers
- **A better solution?**
- For each letter there are 50 numbers
 - So the number of seats is $26 * 50 = 1300$

Product rule: number of letters * number of integers in [1,50]

Product rule

A count can be broken down into a sequence of dependent counts

- “each element in the first count is associated with all elements of the second count”
- **Product rule:** If a count of elements can be broken down into a sequence of dependent counts where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, by the product rule the total number of elements is:
 - $n = n_1 * n_2 * \dots * n_k$


Product rule

Example:

- How many different bit strings of length 7 are there?
 - E.g. 1011010
- Is it possible to decompose the count problem and if yes how?

Yes.

- Count the number of possible assignments to bit 1
- For the first bit assignment (say 0) count assignments to bit 2



Total assignments to first 2 bits: $2*2=4$

Product rule

Example:

- How many different bit strings of length 7 are there?
 - E.g. 1011010
- Is it possible to decompose the count problem and if yes how?
- **Yes.**
 - Count the number of possible assignments to bit 1
 - For the specific first bit count possible assignments to bit 2
 - For the specific first two bits count assignments to bit 3
 - Number of assignments to the first 3 bits: $2*2*2=8$

Product rule

Example:

- How many different bit strings of length 7 are there?
 - E.g. 1011010
- Is it possible to decompose the count problem and if yes how?
- **Yes.**
 - Count the number of possible assignments to bit 1
 - For the specific first bit count possible assignments to bit 2
 - For the specific first two bits count assignments to bit 3
 - Gives a sequence of n dependent counts and by the product rule we have:

$$n = 2 * 2 * 2 * 2 * 2 * 2 * 2 = 2^7$$

Product rule

Example:

The number of subsets of a set S with k elements.

- How to count them?
- **Hint:** think in terms of bit-string representation of a set?
- Assume each element in S is assigned a bit position.
- If A is a subset it can be encoded as a bit-string:
 - if an element is in A then use 1 else put 0
- How many different bit-strings are there?
 - $n = 2 * 2 * \dots 2 = 2$

k bits

Sum rule

A count decomposes into a set of independent counts

- “elements of counts are alternatives”, they do not depend on each other

Example:

- You need to travel in between city A and B. You can either fly, take a train, or a bus. There are 12 different flights in between A and B, 5 different trains and 10 buses. How many options do you have to get from A to B?
- We can take only one type of transportation and for each only one option. The number of options:
 - $n = 12 + 5 + 10$

Sum rule:

- $n = \text{number of flights} + \text{number of trains} + \text{number of buses}$

Sum rule

A count decomposes into a set of independent counts

- “elements of counts are alternatives”
- **Sum rule:** If a count of elements can be broken down into a set of independent counts where the first count yields n_1 elements, the second n_2 elements, and k th count n_k elements, by the sum rule the total number of elements is:
 - $n = n_1 + n_2 + \dots + n_k$

Beyond basic counting rules

- **More complex counting problems** typically require a combination of the sum and product rules.

Example: A login password:

- The minimum password length is 6 and the maximum is 8. The password can consist of either an uppercase letter or a digit. There must be at least one digit in the password.
- How many different passwords are there?

Beyond basic counting rules

Step 1:

- The password we select has either 6,7 or 8 characters.
- So the total number of valid passwords is by the sum rule:

- $P = P_6 + P_7 + P_8$

The number of passwords of length 6,7 and 8 respectively

Step 2:

- Assume passwords with 6 characters (upper-case letters):
- How many are there?
- If we let each character to be at any position we have:

$P_{6\text{-nodigits}} = 26^6$ different passwords of length 6

$P_{6\text{-all}} = (26 + 10)^6 = (36^6)$ different passwords of length 6

Beyond basic counting rules

Step 2

But we must have a password with at least one digit. How to account for it?

A trick. Split the count of all passwords of length 6 into two mutually exclusive groups:

- **$P6\text{-all} = P6\text{-digits} + P6\text{-nodigits}$**
 1. $P6\text{-digits}$ – count when the password has one or more digits
 2. $P6\text{-nodigits}$ – count when the password has no digits
- We know how to easily compute $P6\text{-all}$ and $P6\text{-nodigits}$
 - **$P6\text{-all} = 36^6$ and $P6\text{-nodigits} = 26^6$**
 - Then **$P6\text{-digits} = P6\text{-all} - P6\text{-nodigits}$**

Beyond basic counting rules

Step 1:

the total number of valid passwords is by the sum rule:

- **$P = P6 + P7 + P8$**
- The number of passwords of length 6, 7 and 8 respectively

Step 2

The number of valid passwords of length 6:

$$\begin{aligned} P6 &= P6\text{-digits} = P6\text{-all} - P6\text{-nodigits} \\ &= 36^6 - 26^6 \end{aligned}$$

Analogically:

$$\begin{aligned} P7 &= P7\text{-digits} = P7\text{-all} - P7\text{-nodigits} \\ &= 36^7 - 26^7 \end{aligned}$$

$$\begin{aligned} P8 &= P8\text{-digits} = P8\text{-all} - P8\text{-nodigits} \\ &= 36^8 - 26^8 \end{aligned}$$

Inclusion-Exclusion principle


Used in counts where the decomposition yields two dependent count tasks with overlapping elements

- If we used the sum rule some elements would be counted twice

Inclusion-exclusion principle: uses a sum rule and then corrects for the overlapping elements.

We used the principle for the cardinality of the set union.

- $|A \cup B| = |A| + |B| - |A \cap B|$



Inclusion-Exclusion principle

Example: How many bitstrings of length 8 start either with a bit 1 or end with 00?

- It is easy to count **strings that start with 1**:
- How many are there? 2^7
- It is easy to count the **strings that end with 00**.
- How many are there? 2^6
- Is it OK to add the two numbers to get the answer? $2^7 + 2^6$
- **No. Overcount.** There are some strings that can both start with 1 and end with 00. These strings are counted in twice.
- How to deal with it? How to correct for overlap?
- How many of strings were counted twice? 2^5 (1 xxxxx 00)
- Thus we can correct for the overlap simply by using:
- $2^7 + 2^6 - 2^5 = 128 + 64 - 32 = \mathbf{160}$

Pigeonhole principle

- Assume you have a set of objects and a set of bins/box/basket used to store objects.
- **The pigeonhole principle** states that if there are more objects than bins, then there is at least one bin with more than one object.



n (pigeon) = 5
 m (pigeon hole) = 4



Pigeonhole principle



n (pigeon) = 5

m (pigeon hole) = 4



- **Example:** 5 pigeons and 4 holes
- At least one hole with more than 1 pigeon.

Pigeonhole principle

- Assume you have a set of objects and a set of bins used to store objects.

The pigeonhole principle states that if there are more objects than bins then there is at least one bin with more than one object.

- **Theorem.** If there are $k+1$ objects and k bins. Then there is at least one bin with two or more objects.



K bins

Pigeonhole principle

- Assume you have a set of objects and a set of bins used to store objects. The pigeonhole principle states that if there are more objects than bins then there is at least one bin with more than one object.
- **Theorem.** If there are $k+1$ objects and k bins. Then there is at least one bin with two or more objects.

- **Proof. (by contradiction)**

- Assume that we have $k + 1$ objects and every bin has at most one element. Then the total number of elements is k which is a contradiction.
- End of proof

Pigeonhole principle

Example:

- Assume 367 people. Are there any two people who has the same birthday?
- How many days are in the year? 365.
- Then there must be **at least two people with the same birthday.**

Generalized pigeonhole principle

- We can often say more about the number of objects.
- Say we have 5 bins and 12 objects. What is it we can say about the bins and number of elements they hold?
- There must be **a bin with at least 3 elements**.
- Why?
- Assume there is no bin with more than 3 elements. Then the max number of elements we can have in 5 bins is 10. We need to place 12 so at least one bin should have at least 3 elements.

Generalized pigeonhole principle

Theorem. If N objects are placed into k bins then there is at least one bin containing at least $\lceil N/k \rceil$ objects.

Example. Assume 100 people. Can you tell something about the number of people born in the same month.

- Yes. There exists a month in which at least $\lceil 100 / 12 \rceil = \lceil 8.3 \rceil = 9$ people were born.

; Y b Y f U `] n Y X d] [Y c b \